Recent B Physics Results from CDF


Robert Harr
Wayne State University

Recent B Physics Results from CDF

Using up to 220 pb⁻¹ of data

- Masses and lifetimes
 - A prelude to CP measurements
- Rare decays
- Observation of the state X(3872)
- Hadronic B decay final states are collected with the displaced track trigger
 - $-B_{\rm s}$ mixing channel
 - Access to CP angles □ and □

B Hadron Masses




© Competitive measurements for B_d and B^+ :


 $M(B_d) = 5280.30 \pm 0.92 \pm 0.96 \text{ MeV}$ and $M(B^+) = 5279.32 \pm 0.68 \pm 0.94 \text{ MeV}$

@ World's best measurements for B_s and \square_B :

 $M(B_s) = 5365.50 \pm 1.29 \pm 0.94 \text{ MeV}$ and $M(\square_B) = 5620.4 \pm 1.6 \pm 1.2 \text{ MeV}$

B Hadron Lifetimes







• Check B_d and B^+ lifetimes against BaBar/Belle measurements.

$$[(B_s) = 1.33 \pm 0.14 \pm 0.02 \text{ ps and } [(B_B) = 1.25 \pm 0.26 \pm 0.10 \text{ ps}]$$

- Competitive with LEP results. Will improve with statistics.
- See talk by Daria Zieminska.


Rare Decays






- **Q** New limit for B.R. of $B_s \square \square^{+\square^{-1}}$
 - < 9.5 x 10⁻⁷ @ 90% CL
 - < 1.2 x 10⁻⁶ @ 95% CL
- Limit on B.R. of B_d □ □+□ about a factor of 2 worse than BaBar and Belle.
- New limit for B.R. of
 D⁰ □ □+□-:
 - < 2.5 X 10⁻⁶ @ 90% CL
- See talk by Cheng-Ju Lin.

Hunting For New States



• Ssssshhhh.....


Hunting For New States




• Ssssshhhh......Wabbit hunting.


X(3872)

- New state observed by Belle in $B \square XK \square J/\square \square^+\square^-K$ final state of narrow width.
- ◆ Tevatron: this state produced directly, or via B decays.
- CDF observes this state at the same mass.
- ◆ Belle reports that M(□□) distribution suggests a □ resonance.
- ◆ CDF sees a preference for M(□□) > 500MeV □ needs to be finalized!





X(3872) Signal Significance




- © Note relatively large cross section (times branching fraction) compared to the $\square(2s)$.


What is X(3872)?


- Two leading candidates:
 - 1. A ccbar state \square like the 1 ${}^{3}D_{2}$ state
 - 2. D*D molecule (suggested by Belle)
 - Observed mass is a few MeV below threshold
 - \bullet X \square \square c is not yet observed by Belle
 - \bullet X \square J/ \square \square forbidden for 3D_2 state
- Q Additional measurements to pin down the quantum numbers:
 - Helicity angles
 - \bullet $M_{\square\square}$ distribution (resonance structure)

B□ h+h'- Decays

- $@B_d \square \square \square$ and $B_s \square KK$ modes are sensitive to CP angle \square
- $\bigcirc B_{(d,s)} \square K \square$ and above modes are separated statistically by kinematics and particle ID.
- ② $\mathfrak{B}(B_{d} \square \square)/\mathfrak{B}(B_{d} \square K\square) = 0.26 \pm 0.11 \pm 0.055.$
- **@** First observation of $B_s \square K^+K^-$









- MC templates used to fit mass spectrum.
- $\bigcirc B \sqcap DK$ is another mode of interest for CP analysis.

$B_{\mathrm{S}} \square D_{\mathrm{S}}^{-} \square^{+}$





- $\bigcirc B_{\rm s} \square D_{\rm s}$ is a potential mode for measuring $x_{\rm s}$.
- © Fully reconstructed mode for minimal uncertainty in the boost of the B_s .

Summary

- Progress with CDF Run2 analyses:
 - **♦** Traditional: masses, lifetimes, rare decays.
 - \bullet CDF has observed the X(3872).
 - New topics enabled by the displaced track trigger are being explored: hadronic CP modes, and hadronic modes for B_s mixing.
- New charm and bottom production cross sectionsChunhui Chen.
- © Tagging studies [Ting Miao.